



Likelihood for Alfvénic bifurcation in experiments

Vinícius Duarte^{1,2}

In collaboration with

Nikolai Gorelenkov¹, Herb Berk³, Eric Fredrickson¹, Mia Schneller¹, Mario Podestà¹, Raffi Nazikian¹,
Mike Van Zeeland⁴, David Pace⁴ and Bill Heidbrink⁵

¹Princeton Plasma Physics Laboratory, USA

²University of São Paulo, Brazil

³University of Texas, Austin, USA

⁴General Atomics, USA

⁵University of California, Irvine, USA

IAEA Energetic Particle Meeting – Princeton, September 5th 2017

Outline

- Introduction on Alfvénic spectral characteristics induced by energetic particles
- The cubic equation (Berk-Breizman model) and a criterion for onset of chirping
- Micro-turbulence as a mediator for chirping onset in DIII-D and NSTX
- Predictions for ITER


Outline

- **Introduction on Alfvénic spectral characteristics induced by energetic particles**
- The cubic equation (Berk-Breizman model) and a criterion for onset of chirping
- Micro-turbulence as a mediator for chirping onset in DIII-D and NSTX
- Predictions for ITER


Alfvén waves can exhibit a range of bifurcations upon their interaction with fast ions

Typical scenarios:


- fixed frequency and frequency splitting-> frequency is mostly determined by the equilibrium
- chirping and avalanches -> frequency is highly affected by the fast ions nonlinear response



Fasoli, PRL 1998



Fredrickson, PoP 2006



Podestà, NF 2011

Prediction of character of energetic-particle-driven transport in tokamaks

What tools can be used to model each type of transport?

Diffusive transport (typical for fixed-frequency modes)

- can be modelled using reduced theories, such as quasilinear
- typical in conventional tokamaks

Convective transport (typical for chirping frequency modes)

- needs to retain full nonlinear features of the wave, is sustained by nonlinear phase-space structures
- typical in spherical tokamaks

Prediction of character of energetic-particle-driven transport in tokamaks


What tools can be used to model each type of transport?

Diffusive transport (typical for fixed-frequency modes)

- can be modelled using reduced theories, such as quasilinear
- typical in conventional tokamaks

Convective transport (typical for chirping frequency modes)

- needs to retain full nonlinear features of the wave, is sustained by nonlinear phase-space structures
- typical in spherical tokamaks



Both can lead to similar fast ion loss levels, up to 40% in present-day experiments

Prediction of character of energetic-particle-driven transport in tokamaks


What tools can be used to model each type of transport?

Diffusive transport (typical for fixed-frequency modes)

- can be modelled using reduced theories, such as quasilinear
- typical in conventional tokamaks

Convective transport (typical for chirping frequency modes)

- needs to retain full nonlinear features of the wave, is sustained by nonlinear phase-space structures
- typical in spherical tokamaks



Both can lead to similar fast ion loss levels, up to 40% in present-day experiments

In this talk:

Prediction of character of energetic-particle-driven transport in tokamaks


What tools can be used to model each type of transport?

Diffusive transport (typical for fixed-frequency modes)

- can be modelled using reduced theories, such as quasilinear
- typical in conventional tokamaks

Convective transport (typical for chirping frequency modes)

- needs to retain full nonlinear features of the wave, is sustained by nonlinear phase-space structures
- typical in spherical tokamaks



Both can lead to similar fast ion loss levels, up to 40% in present-day experiments

In this talk:

- **development of a criterion for the likelihood of each nonlinear scenario and its comparison with NSTX and DIII-D**

Prediction of character of energetic-particle-driven transport in tokamaks


What tools can be used to model each type of transport?

Diffusive transport (typical for fixed-frequency modes)

- can be modelled using reduced theories, such as quasilinear
- typical in conventional tokamaks

Convective transport (typical for chirping frequency modes)

- needs to retain full nonlinear features of the wave, is sustained by nonlinear phase-space structures
- typical in spherical tokamaks




Both can lead to similar fast ion loss levels, up to 40% in present-day experiments

In this talk:

- development of a criterion for the likelihood of each nonlinear scenario and its comparison with NSTX and DIII-D
- Predictions for TAE in ITER elmy and hybrid scenarios


The study of the conditions that lead to fully nonlinear scenarios helps to understand the applicability of reduced models

Need for predictive/efficient
interpretive capabilities motivates
the development of phase-space
resolved, self-consistent quasilinear
approach




The study of the conditions that lead to fully nonlinear scenarios helps to understand the applicability of reduced models

Need for predictive/efficient interpretive capabilities motivates the development of phase-space resolved, self-consistent quasilinear approach




Diffusive module of the Resonance Broadening Quasilinear (RBQ) code, for the case of two overlapping resonances




[Detailed description of the RBQ code in N. Gorelenkov's poster]

The study of the conditions that lead to fully nonlinear scenarios helps to understand the applicability of reduced models

Need for predictive/efficient interpretive capabilities motivates the development of phase-space resolved, self-consistent quasilinear approach




Diffusive module of the Resonance Broadening Quasilinear (RBQ) code, for the case of two overlapping resonances



[Detailed description of the RBQ code in N. Gorelenkov's poster]

DIII-D discharge 153072



[Resonance broadening parametric dependencies in G. Meng's poster] 4

Outline

- Introduction on Alfvénic spectral characteristics induced by energetic particles
- **The cubic equation (Berk-Breizman model) and a criterion for onset of chirping**
- Micro-turbulence as a mediator for chirping onset in DIII-D and NSTX
- Predictions for ITER

Weak nonlinear dynamics of driven kinetic systems can be used to develop a criterion to distinguish between fixed-frequency and chirping responses

Starting point: kinetic equation plus wave power balance

Assumptions:

- Perturbative procedure for $\omega_b \ll \gamma$ ($\omega_b \propto \sqrt{\text{mode amplitude}}$)
- Truncation at third order due to closeness to marginal stability
- Bump-on-tail modal problem, uniform mode structure

Cubic equation: lowest-order nonlinear correction to the evolution of mode amplitude A :

$$\frac{dA}{dt} = A - \int_0^{t/2} d\tau \tau^2 A(t - \tau) \int_0^{t-2\tau} d\tau_1 e^{-\nu_{scatt}^3 \tau^2 (2\tau/3 + \tau_1) + i\nu_{drag}^2 \tau(\tau + \tau_1)} A(t - \tau - \tau_1) A^*(t - 2\tau - \tau_1)$$

Weak nonlinear dynamics of driven kinetic systems can be used to develop a criterion to distinguish between fixed-frequency and chirping responses


Starting point: **kinetic equation** plus **wave power balance**

Assumptions:

- Perturbative procedure for $\omega_b \ll \gamma$ ($\omega_b \propto \sqrt{\text{mode amplitude}}$)
- Truncation at third order due to closeness to marginal stability
- Bump-on-tail modal problem, uniform mode structure

Cubic equation: lowest-order nonlinear correction to the evolution of mode amplitude A :

$$\frac{dA}{dt} = A - \int_0^{t/2} d\tau \tau^2 A(t - \tau) \int_0^{t-2\tau} d\tau_1 e^{-\nu_{scatt}^3 \tau^2 (2\tau/3 + \tau_1) + i\nu_{drag}^2 (\tau + \tau_1)} A(t - \tau - \tau_1) A^*(t - 2\tau - \tau_1)$$



stabilizing destabilizing (makes integral sign flip)

Weak nonlinear dynamics of driven kinetic systems can be used to develop a criterion to distinguish between fixed-frequency and chirping responses


Starting point: kinetic equation plus wave power balance

Assumptions:

- Perturbative procedure for $\omega_b \ll \gamma$ ($\omega_b \propto \sqrt{\text{mode amplitude}}$)
- Truncation at third order due to closeness to marginal stability
- Bump-on-tail modal problem, uniform mode structure

Cubic equation: lowest-order nonlinear correction to the evolution of mode amplitude A :

$$\frac{dA}{dt} = A - \int_0^{t/2} d\tau \tau^2 A(t - \tau) \int_0^{t-2\tau} d\tau_1 e^{-\nu_{scatt}^3 \tau^2 (2\tau/3 + \tau_1) + i\nu_{drag}^2 (\tau + \tau_1)} A(t - \tau - \tau_1) A^*(t - 2\tau - \tau_1)$$



stabilizing destabilizing (makes integral sign flip)

- If nonlinearity is weak: linear stability, solution saturates at a low level and f merely flattens (system not allowed to further evolve nonlinearly).
- If solution of cubic equation explodes: system enters a strong nonlinear phase with large mode amplitude and can be driven unstable (precursor of chirping modes).

A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to


$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int$$
$$Int \equiv Re \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + iz^2 \right]$$

A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int \quad \text{---} \left\{ \begin{array}{l} >0: \text{fixed-frequency solution likely} \\ <0: \text{chirping likely to occur} \end{array} \right.$$

$$Int \equiv Re \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + iz^2 \right]$$




A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int \quad \left\{ \begin{array}{l} >0: \text{fixed-frequency solution likely} \\ <0: \text{chirping likely to occur} \end{array} \right.$$

$$Int \equiv \text{Re} \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + iz^2 \right]$$

Phase space integration



A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to


$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int \quad \left\{ \begin{array}{l} >0: \text{fixed-frequency solution likely} \\ <0: \text{chirping likely to occur} \end{array} \right.$$

$$Int \equiv \text{Re} \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + iz^2 \right]$$

Phase space integration

Eigenstructure information:

$$q \int dt \mathbf{v}_{dr} \cdot \delta \mathbf{E} e^{i\omega t}$$



A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int \quad \left. \begin{array}{l} >0: \text{fixed-frequency solution likely} \\ <0: \text{chirping likely to occur} \end{array} \right\}$$

$$Int \equiv \text{Re} \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + iz^2 \right]$$


Phase space integration

Eigenstructure information:

$$q \int dt \mathbf{v}_{dr} \cdot \delta \mathbf{E} e^{i\omega t}$$

Resonance surfaces:

$$\begin{aligned} \Omega_l(\mathcal{E}' + \omega P_{\varphi}/n, P_{\varphi}, \mu) &\equiv \\ &\equiv n \frac{d\varphi}{dt} - l \frac{d\theta}{dt} - \omega_0 \end{aligned}$$



A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int \quad \left. \begin{array}{l} >0: \text{fixed-frequency solution likely} \\ <0: \text{chirping likely to occur} \end{array} \right\}$$

$$Int \equiv Re \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + i z^2 \right]$$

Phase space integration


Eigenstructure information:

$$q \int dt \mathbf{v}_{dr} \cdot \delta \mathbf{E} e^{i\omega t}$$

Resonance surfaces:

$$\begin{aligned} \Omega_l(\mathcal{E}' + \omega P_{\varphi}/n, P_{\varphi}, \mu) &\equiv \\ &\equiv n \frac{d\varphi}{dt} - l \frac{d\theta}{dt} - \omega_0 \end{aligned}$$

Criterion was incorporated into NOVA-K code:
nonlinear prediction from linear physics elements



A criterion for the likelihood of chirping onset in tokamaks

Using an action and angle formulation, the previous weak nonlinear theory leads to

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \frac{\partial f}{\partial I} Int \quad \left. \begin{array}{l} >0: \text{fixed-frequency solution likely} \\ <0: \text{chirping likely to occur} \end{array} \right\}$$

$$Int \equiv \text{Re} \int_0^{\infty} dz \frac{z}{\frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z - i} \exp \left[-\frac{2}{3} \frac{\nu_{\text{stoch}}^3}{\nu_{\text{drag}}^3} z^3 + i z^2 \right]$$

Phase space integration


Eigenstructure information:

$$q \int dt \mathbf{v}_{dr} \cdot \delta \mathbf{E} e^{i\omega t}$$

Resonance surfaces:

$$\begin{aligned} \Omega_l(\mathcal{E}' + \omega P_{\varphi}/n, P_{\varphi}, \mu) &\equiv \\ &\equiv n \frac{d\varphi}{dt} - l \frac{d\theta}{dt} - \omega_0 \end{aligned}$$

Criterion was incorporated into NOVA-K code:
nonlinear prediction from linear physics elements






The criterion ($Crt \geq 0$) predicts that micro-turbulence should be key in determining the likely nonlinear character of a mode, e.g., fixed-frequency or chirping




Outline

- Introduction on Alfvénic spectral characteristics induced by energetic particles
- The cubic equation (Berk-Breizman model) and a criterion for onset of chirping
- **Micro-turbulence as a mediator for chirping onset in DIII-D and NSTX**
- Predictions for ITER

Correlation between chirping onset and a marked reduction of the turbulent activity in DIII-D




- The thermal ion heat conductivity is used as a proxy for the fast ion anomalous transport
- This observation motivated DIII-D experiments to be designed to further test the hypothesis of low turbulence associated with chirping



Dedicated experiments showed that chirping is more prevalent in negative triangularity DIII-D shots

- Transport coefficients calculated in TRANSP are 2-3 times lower in negative triangularity, as compared to the the usual positive triangularity/oval shots;
- Chirping found to exist in positive triangularity at the bottom of RSAEs evolution, where an ITB is expected.



[Detailed description of the negative triangularity experiments given in M. Van Zeeland's talk this morning]



DIII-D: chirping criterion evaluation in negative vs positive triangularity

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \left| \frac{\partial f}{\partial I} \right|_{\text{Int}}$$



>0: fixed-frequency solution likely
<0: chirping likely to occur





DIII-D: chirping criterion evaluation in negative vs positive triangularity

$$Crt = \frac{1}{N} \sum_{j, \sigma_{\parallel}} \int dP_{\varphi} \int d\mu \frac{|V_j|^4}{\omega_{\theta} \nu_{\text{drag}}^4} \left| \frac{\partial \Omega_j}{\partial I} \right| \left| \frac{\partial f}{\partial I} \right|_{\text{Int}}$$



>0: fixed-frequency solution likely
<0: chirping likely to occur



Correlation between chirping onset and a marked reduction of the turbulent activity in NSTX, as computed by TRANSP





- The thermal ion heat conductivity is used as a proxy for the fast ion anomalous transport






- GTS code is being used to verify earlier publications on the EP turbulence-induced anomalous scattering

GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping




*W.X. Wang et al., PoP **13**, 092505 (2006)


GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping





GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping





GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping





GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping



GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping




GTS* global gyrokinetics analyses show turbulence reduction for rare NSTX TAE transitions from fixed-frequency to chirping




Examples of the chirping criterion evaluation: spherical vs conventional tokamaks

chirping, NSTX

Alfvén wave
chirping
quantitatively
agrees with
the criterion



fixed-frequencies, DIII-D and TFTR



Arrows
represent the
turbulent
diffusion that
adds up to
pitch-angle
scattering


Chirping is ubiquitous in NSTX but rare in DIII-D, which is consistent with the inferred fast ion micro-turbulent levels

Outline

- Introduction on Alfvénic spectral characteristics induced by energetic particles
- The cubic equation (Berk-Breizman model) and a criterion for onset of chirping
- Micro-turbulence as a mediator for chirping onset in DIII-D and NSTX
- **Predictions for ITER**

Predictions for n=7-11 TAEs¹ in ITER are near threshold between steady and chirping

Based on TRANSP/TSC analysis, requiring $Q > 10$



Approximate rate of radial turbulence diffusion to the collisional pitch-angle scattering^{2,3}

$$\text{Ratio} \approx \frac{D_{EP} \left(\frac{q_{EP}}{m_{EP}} \frac{\partial \psi}{\partial r} \right)^2}{2\nu_{\perp} R^2 \left[\mathcal{E} - \frac{B_{\varphi}^2}{B^2} (\mathcal{E} - \mu B) \right]}$$

$$\nu_{\perp} = \frac{1}{2} \langle Z \rangle \frac{\bar{A}_i}{[Z]} \frac{1}{A_{EP}} \left(\frac{v_c}{v} \right)^3 \frac{1}{\tau_s}$$

¹ DOE OFES Theory Joule Milestone FY2007, Gorelenkov et al, PPPL Preprint number 4287 (2008).

²Lang & Fu, PoP 2010.

³Duarte et al, NF 2017.

Summary

- Criterion gives confidence in the application of quasilinear modeling;
- The gyrokinetic code GTS confirms transition from/to chirping is likely mediated by a change of turbulence;
- Experiments with negative triangularity on DIII-D give credence to the proposed chirping criterion predictions;
- Predicted response for ITER (similarly to DIII-D predictions) appears to be around the borderline between fixed-frequency and chirping.

Thank you